


FACULTY OF NURSING



UNIT-10 OCCUPATIONAL AND INDUSTRIAL HEALTH DISORDER

BY- ANWAR SHAHJAD M.SC.(N) LECTURER



OCCUPATIONAL HEALTH

WHO DEFINED OCCUPATIONAL HEALTH AS;

'THE PROMOTION AND MAINTENANCE OF THE HIGHEST DEGREE OF PHYSICAL, MENTAL AND SOCIAL WELL-BEING OF WORKERS IN ALL OCCUPATIONS



WHAT IS OCCUPATIONAL HEALTH NURSING?

>APPLICATION OF NURSING PRINCIPLES IN CONSERVING THE HEALTH OF WORKERS AT THE WORKPLACE.

> BASED ON PRINCIPLES OF RECOGNITION, PREVENTION AND TREATMENT OF ILLNESS, INJURIES.

FORMALLY KNOWN AS 'INDUSTRIAL NURSING'

AIMS OF OCCUPATIONAL HEALTH NURSING PROMOTION AND MAINTENANCE OF THE HIGHEST **DEGREE OF PHYSICAL, MENTAL AND SOCIAL WELL-BEING OF WORKERS IN ALL OCCUPATIONS ;** BY PREVENTING- DEPARTURES FROM HEALTH **CAUSED BY THEIR WORKING CONDITIONS,** PROTECTING THE WORKERS IN THEIR EMPLOYMENT FROM RISKS RESULTING FROM FACTORS ADVERSE TO **HEALTH AND ADAPTATION OF WORKERS TO THEIR** JOBS

OBJECTIVES OF OCCPATIONAL HEALTH

NURSING

MAINTENANCE AND PROMOTION OF WORKERS' HEALTH AND WORKING CAPACITY(HEALTH MONITORING)

IMPROVEMENT OF WORKING ENVIRONMENT AND WORK TO BECOME CONDUCIVE TO SAFETY AND HEALTH (WORKSITE VISITS)

DEVELOPMENT OF WORK ORGANISATIONS AND WORKING CULTURES IN A DIRECTION, WHICH SUPPORTS HEALTH AND SAFETY AT WORK(HEALTH EDUCATION)

OCCPATIONAL HAZARDS

- **1. PHYSICAL HAZARDS**
- 2. CHEMICAL
- 3. **BIOLOGICAL**
- 4. PSYCHOSOCIAL a.PSYCHOLOGICAL AND BEHAVIORAL CHANGES. b.PSYCHOSOMATIC ILL-HEALTH



1. PHYSICAL HAZARDS

HEAT- DIRECT & INDIRECT EFFECT OF HIGH TEMPERATURE,

RADIANT HEAT, HEAT STAGNATION.

COLD-GENERAL & LOCAL COLD INJURY

LIGHT – ACUTE & CHRONIC EFFECT OF BRIGHT & DIM LIGHT & GLARE.

- RADIATION
 - a. IONIZING-X-RAYS, GAMMARAYS, BETA PARTICLES, ALPHA PARTICLES
 - b. NON-IONIZING-MICROWAVES, INFRARED, ANDULTRA-VIOLET LIGHT

NOISE – AUDITORY & NON AUDITORY EFFECT.

VIBRATION - HAZARDOUS IN THE FREQUENCY RANGE OF 10-500HZ

DISEASES DUE TO PHYSICAL AGENTS

HEAT - HEAT STROKE, HEAT RASH.

LIGHT - OCCUPATIONAL CATARACT

COLD-HYPOTHERMIA, FROST BITE

VIBRATION – OSTEOARTHRITIS

NOISE - OCCUPATIONAL DEAFNESS

RADIATION- CANCER

Frostbite BULLAE

Exposure to a temperature below freezing leads to severe soft tissue damage.

FINGERS WITH FROSTBITE

silebra

- 2. CHEMICAL HAZARDS:-
- ACIDS
- BASES
- HEAVY METALS -LEAD
- SOLVENTS PETROLEUM
- PARTICULATES ASBESTOS, SILICA AND OTHER FINE DUST/FIBROUS MATERIALS
- FUMES NOXIOUS GASES/VAPORS

HIGHLY-REACTIVE METALS

DISEASES DUE TO CHEMICAL AGENTS

- GASES GAS POISONING.
- INORGANIC DUSTS:
 - > COAL DUST ANTHRACOSIS
 - > SILICA SILICOSIS
 - > ASBESTOS ASBESTOSIS
 - > IRON SIDEROSIS
- ORGANIC DUSTS:
 - > CANE FIBER BAGASSOSIS.
 - > COTTON DUST BYSSINOSIS.
 - > HAY OR GRAIN DUST FARMER'S LUNG.
- CHEMICALS BURNS, DERMATITIS, CANCER, RESPIRATORY ILLNESS
- METALS LEAD, MERCURY, ARSENIC, CHROMIUM CAUSE POISONING

PNEUMOCONIOSES:-

IT IS A OCCUPATIONAL LUNG DISEASE CAUSED BY

INHALATION OF DUST.

TYPES:-

- **O COAL WORKER PNEUMOCONIOSES**
- **OBASBESTOSIS**
- **OSILICOSIS**
- BAUXITE FIBROSIS-IT IS A FORM OF
 PNEUMOCONIOSES CAUSED BY BAUXITE
- **BERYLLIOSIS-BERYLLIUM**
- **O SIDEROSIS-IRON**
- **BYSSINOSIS** -COTTON

COAL WORKERS' PNEUMOCONIOSIS - BLACK LUNG DISEASE - EXPOSURE TO PARTICLES OF CARBON


WHEN COAL DUST IS INHALED FOR A LONG PERIOD OF TIME, IT BUILDS UP IN THE LUNGS, WHICH THE BODY IS NOT ABLE TO REMOVE - INFLAMMATION OF THE LUNGS - FIBROSIS - CAUSES LARGE SIZE CAVITIES IN THE LUNGS.



Healthy Tissue 90-year-old schoolteacher Progressive massive fibrosis 40-year-old-miner

SILICOSIS:- LUNG FIBROSIS CAUSED BY THE

INHALATION OF DUST CONTAINING SILICA



BERYLLIOSIS OR CHRONIC BERYLLIUM DISEASE (CBD):-

IT IS A CHRONIC ALLERGIC-TYPE LUNG RESPONSE AND CHRONIC LUNG DISEASE CAUSED BY EXPOSURE TO BERYLLIUM AND ITS COMPOUNDS, A FORM OF BERYLLIUM POISONING

SIDEROSIS (WELDER'S DISEASE):- IT IS THE DEPOSITION OF EXCESS IRON IN BODY TISSUE



CAUSED BY THE ACCUMULATION OF CARBON IN THE LUNGS DUE TO REPEATED EXPOSURE TO AIR POLLUTION OR INHALATION OF COAL DUST PARTICLES"

ANTHRACOSIS :- IT'S A TYPE OF PNEUMOCONIOSIS AS

FARMER'S LUNG IS A DISEASE CAUSED BY AN ALLERGY TO THE CERTAIN CROPS. FARMERS ARE MOST LIKELY TO GET IT BECAUSE IT'S USUALLY CAUSED BY BREATHING IN DUST FROM HAY, CORN, GRASS FOR ANIMAL FEED, GRAIN, TOBACCO, OR SOME PESTICIDES.

ASBESTOSIS:-

ASBESTOSIS (AS-BES-TOE-SIS) IS A CHRONIC LUNG DISEASE CAUSED BY INHALING ASBESTOS FIBERS.

PROLONGED EXPOSURE TO THESE FIBERS CAN CAUSE LUNG TISSUE SCARRING AND SHORTNESS OF BREATH.

ASBESTOSIS SYMPTOMS CAN RANGE FROM MILD TO SEVERE, AND USUALLY DON'T APPEAR UNTIL MANY YEARS AFTER CONTINUED EXPOSURE.

ASBESTOS IS A NATURAL MINERAL PRODUCT THAT'S RESISTANT TO HEAT AND CORROSION. IT WAS USED IN PRODUCTS SUCH AS CEMENT AND SOME FLOOR TILES.

REDUCING EXPOSURE TO ASBESTOS IS THE BEST PREVENTION AGAINST ASBESTOSIS

Bagassosis

 Inhalation of bagasse sugar cane dust containing thermophilic actinomycete, thermoactinomyces sacchari

Symptoms:

Breathlessness, cough, haemoptysis, and slight fever.

Occupational exposure:

Manufacturing of paper, cardboard and rayon.



Byssinosis

- Byssinosis:
 - Caused by inhalation of cotton fibre dust
 - The chief symptoms are
 - Chest tightness
 - · Shortness of breath
 - · Cough and
 - Wheezing



- Typically occurring when patients return to work after a weekend or vacation.
- Smoking significantly exacerbates byssinosis

3. BIOLOGICAL HAZARDS

BACTERIA VIRUSES

FUNGI

INSECTS – MOSQUITOES

ANIMALS



DISEASES DUE TO BIOLOGICAL AGENTS

HEPATITIS B VIRUS

HEPATITIS C VIRUS

TUBERCULOSIS



4. PSYCHOSOCIAL HAZARDS WORK-RELATED STRESS – EXCESSIVE WORKING TIME AND OVERWORK

VIOLENCE – FROM OUTSIDE THE ORGANIZATION BULLYING – EMOTIONAL AND VERBAL ABUSE SEXUAL HARASSMENT

MOBBING

BURNOUT

EXPOSURE TO UNHEALTHY ELEMENTS – TOBACCO,

UNCONTROLLED ALCOHOL

DISORDERS DUE TO PSYCHOSOCIAL HAZARDS MUSCULOSKELETAL DISORDERS (MSDS) •CAN AFFECT THE BODY'S MUSCLES, JOINTS, **TENDONS, LIGAMENTS AND NERVES. AS WELL AS** THE BACK, NECK, SHOULDERS AND UPPER LIMBS; LESS OFTEN THEY AFFECT THE LOWER LIMBS. **OMEMORY LOSS DISSATISFACTION OFRUSTRATION & IRRITABILITY**

- MEDICAE^{NTIO}METHODS^S SUCH AS PERIODIC EXAMINATIONS, PREPLACEMENT, WORKING ENVIRONMENT SUPERVISION AND HEALTH EDUCATION. NOTIFICATION, MAINTENANCE AND ANALYSIS OF RECORDS, AND COUNSELLING ARE ALSO STEPS TOWARDS PREVENTION.
- ENGINEERING MEASURES SUCH AS PROPER DESIGN OF PLANT, VENTILATION, DUST ISOLATION THROUGH ENCLOSURE AND ISOLATION, AND PROTECTIVE DEVICES.
- LEGISLATIVE PROTECTION LIKE THE FACTORIES ACT OF 1948 AND THE EMPLOYEES STATE INSURANCE ACT OF 1948.

PREVENTION OF OCCUPATIONAL DISEASES MEASURES FOR HEALTH PROTECTION OF WORKERS.

- 1. NUTRITION
- 2. COMMUNICABLE DISEASE CONTROL
- 3. ENVIRONMENTAL SANITATION.
 - WATER SUPPLY
 - FOOD
 - TOILET
 - PROPER GARBAGE & WASTE DISPOSAL.
 - GENERAL PLANT CLEANLINESS.
 - SUFFICIENT SPACE.
 - LIGHTING.
 - VENTILATION.
 - PROTECTION AGAINST HAZARDS.

4. MENTAL HEALTH:

GOALS:

TO PROMOTE HEALTH AND HAPPINESS OF WORKERS.

TO DETECT SIGNS OF STRESS AND STRAIN AND TAKE NECESSARY MEASURES.

TREATMENT OF EMPLOYEES SUFFERING FROM

MENTAL ILLNESSES.

REHABILITATION OF THOSE WHO BECOME ILL

5. HEALTH EDUCATION.



PREVENTION OF OCCUPATIONAL DISEASE

PRIMARY PREVENTION IS DESIGNED TO DETER OR AVOID THE OCCURRENCE OF DISEASE OR INJURY

SECONDARY PREVENTION IS DESIGNED TO IDENTIFY AND ADEQUATELY TREAT A DISEASE OR INJURY PROCESS AS SOON AS POSSIBLE, OFTEN BEFORE ANY SYMPTOMS HAVE DEVELOPED

TERTIARY PREVENTIONIS DESIGNED TO TREAT A DISORDER
WHEN IT HAS ADVANCED BEYOND
ITS EARLY STAGES, TO AVOID
COMPLICATIONS & LIMIT
DISABILITY, TO ADDRESS
REHABILITATIVE AND PALLIATIVE
NEEDS

