



**FACULTY OF AGRICULTURAL SCIENCES AND
ALLIED INDUSTRIES**

AGR- 121 Soil and Water Conservation Engineering

Lecture 1

Soil Erosion: Introduction

What is Soil Erosion?

The uppermost weathered and disintegrated layer of the earth's crust is referred to as soil. The soil layer is composed of mineral and organic matter and is capable of sustaining plant life. The soil depth is less in some places and more at other places and may vary from practically nil to several metres. The soil layer is continuously exposed to the actions of atmosphere. Wind and water in motion are two main agencies which act on the soil layer and dislodge the soil particles and transport them. The loosening of the soil from its place and its transportation from one place to another is known as soil erosion.

The word erosion has been derived from the Latin word 'erodere' which means eating away or to excavate. The word erosion was first used in geology for describing the term hollow created by water. Erosion actually is a two phase process involving the detachment of individual soil particle from soil mass, transporting it from one place to another (by the action of any one of the agents of erosion, viz; water, wind, ice or gravity) and its deposition. When sufficient energy is not available to transport a particle, a third phase known as deposition occurs. In general, finer soil particles get eroded more easily than coarse particles (silt is more easily eroded than sand). Hence soil erosion is defined as a process of detachment, transportation and deposition of soil particles (sediment). It is evident that sediment is the end product of soil erosion process. Sediment is, therefore, defined as any fragmented material, which is transported or deposited by water, ice, air or any other natural agent. From this, it is inferred that sedimentation is also the process of detachment, transportation and deposition of eroded soil particles. Thus, the natural sequence of the sedimentation is as follows:

SOIL DETACHMENT → **TRANSPORT** → **DEPOSITION**

Detachment is the dislodging of the soil particle from the soil mass by erosive agents. In case of water erosion, major erosive agents are impacting raindrops and runoff water flowing over the soil surface. Transportation is the entrainment and movement of detached soil particles (sediment) from their original location. Sediments move from the upland sources through the stream system and may eventually reach the ocean. Not all the sediment reaches the ocean; some are deposited at the base of the slopes, in reservoirs and flood plains along the way. Erosion is almost universally recognized as a serious threat to human

well being. Erosion reduces the productivity of crop land by removing and washing away of plant nutrients and organic matter.

Balanced ecosystems comprising soil, water and plant environments are essential for the survival and welfare of mankind. However, ecosystems have been disturbed in the past due to over exploitation in many parts of the world, including some parts of India. The resulting imbalance in the ecosystem is revealed through various undesirable effects, such as degradation of soil surfaces, frequent occurrence of intense floods etc. Vast tracts of land have been irreversibly converted into infertile surfaces due to accelerated soil erosion caused by the above and other factors. These degraded land surfaces have also become a source of pollution of the natural water. Deposition of soil eroded from upland areas in the downstream reaches of rivers has caused aggradation. This has resulted in an increase in the flood plain area of the rivers, reduction of the clearance below bridges and culverts and sedimentation of reservoirs.

The major land degradation problems due to sedimentation are briefly discussed as below:

- **Erosion by wind and water:** Out of 144.12 M-ha areas affected by water and wind erosion. About 69 M-ha is considered to be critical and needs immediate attention. Wind erosion is mainly restricted to States of Rajasthan, Gujarat and Haryana. The severity of wind erosion is inversely related to the rainfall amount, lesser is the rainfall more would be the wind erosion.
- **Gullies and Ravines:** About 4 M-ha is affected by the problem of gullies and ravines in the country covering about 12 states. Ravines are mostly located in the states of Uttar Pradesh, Madhya Pradesh, Rajasthan and Gujarat. Gullies on the other hand are seen in the plateau region of Eastern India, foot hills of the Himalayas and areas of Deccan Plateau.
- **Torrents and Riverine Lands:** Problem of Riverine and torrents is spread over an area of 2.73 M-ha in the country. Torrents are the natural streams which cause extensive damage to life and property as a result of frequent changes in their course and associated flash flows with heavy debris loads. The unfertile material or debris transported by torrents is sometimes deposited on the fertile plains, thus ruining the land for ever.
- **Water logging:** Water logging is caused either by surface flooding or due to rise of water table. An area of 8.53 M-ha has been estimated to be affected by water logging. Water logging due to surface flooding is predominant in the states of West

Bengal, Assam, Bihar, Orissa, Andhra Pradesh, Uttar Pradesh, Kerala, Punjab and Haryana.

- **Shifting Cultivation:** Shifting cultivation, also known as 'jhuming' is a traditional method of growing crops on hill slopes by slash and burn method. The method involves selection of appropriate site on hill slopes, cleaning of forest by cutting and burning, using the site for cultivation for few years and later on abandoning it and moving to a fresh site. The jhum cycle has gradually declined from 20-30 years to 3-6 years due to increasing population pressures. The problem is more serious in North Eastern region and in the states of Orissa and Andhra Pradesh.
- **Saline soil including coastal areas:** Saline soils are prevalent both in inland as well as coastal areas. About 5.5 M-ha area is affected by this problem in the country which includes arid and semi-arid areas of Rajasthan and Gujarat, black soil region and coastal areas. This problem is causing serious damage to agricultural lands, rendering fertile soil unproductive and turning groundwater brackish in the States of West Bengal, Tamil Nadu, Orissa, Maharashtra, Kerala, Karnataka, Gujarat and Andhra Pradesh as well as Union Territories of Pondicherry and Goa, Daman and Diu.
- **Floods and Droughts:** In India, among the major and medium rivers of both Himalayas and non-Himalayas categories, 18 are flood prone which drain an area of 150 M-ha. In recent years, flash floods have caused extensive damage even in the desert areas of Rajasthan and Gujarat.

1.1.3 Importance of Soil Conservation

In India, out of the total geographical area of 329 M-ha, an area of about 150 M-ha is subjected to either water or wind erosion. A net area of about 140 M-ha is cropped at present. An area of 40 M-ha is considered to be flood prone. Area lost through ravines and gullies is estimated to be about 4 M-ha. As a whole, it is estimated that about 175 M-ha i.e., 53.3% of the total geographical area of the country is subjected to various soil and land degradation problems like saline-alkali soils, waterlogged areas, ravine and gullied lands, area under shifting cultivation, and desertification. By the year 2100 A.D, the projected population of the country is expected to be two billion, whereas the food grain production is almost stagnant at 211 million tons for the last 5 years. The per capita cropped area is shrinking every day; in the year 1950, it was 0.33 ha/capita, 0.2 ha in 1980 and it was 0.15 ha by 2000. This clearly shows that the limited land resource has to be managed very carefully by adopting total conservation measures for the survival of the huge population. A few suggestions to conserve soil and water resources in Indian context are discussed below.

- To prevent erosion of bare soil, it is important to maintain a vegetation cover, especially in the most vulnerable areas e.g. those with steep slopes, in a dry season or periods of very heavy rainfall. For this purpose, only partial harvesting forests (e.g.

alternate trees) and use of seasonally dry or wet areas for pasture rather than arable agricultural land should be permitted.

- Where intensive cultivation takes place, farmers should follow crop rotation in order to prevent the soil becoming exhausted of organic matters and other soil building agents. Where soils are ploughed in vulnerable areas, contour ploughing (i.e. round the hillside rather than down the hillside) should be used. Careful management of irrigation, to prevent the application of too much or too little water will be helpful to reduce the problem of soil salinity development. Livestock grazing must be carefully managed to prevent overgrazing.
- Construction of highways and urbanization should be restricted to areas of lower agricultural potential. With extractive industries, a pledge must be secured to restore the land to its former condition before permission for quarries or mines is granted.