ENVIRONMENTAL STUDIES

LECTURE-18

Ecological succession

Succession is the order of colonization of species in an ecosystem from a barren or destroyed area of land. Mosses and lichens are the first species that inhabit an area. They make the area suitable for the growth of larger species such as grasses, shrubs and finally trees.

"Ecological succession is a series of changes that occur in an ecological community over time."

Ecological succession is the steady and gradual change in a species of a given area with respect to the changing environment. It is a predictable change and is an inevitable process of nature as all the biotic components have to keep up with the changes in **our environment**.

The ultimate aim of this process is to reach equilibrium in the ecosystem. The community that achieves this aim is called a climax community. In an attempt to reach this equilibrium, some species increase in number while some other decrease.

In an area, the sequence of communities that undergo changes is called sere. Thus, each community that changes is called a seral stage or seral community.

All the communities that we observe today around us have undergone succession over a period of time since their existence. Thus, we can say that evolution is a process that has taken place simultaneously along with that of ecological succession. Also, the initiation of life on earth can be considered to be a result of this succession process.

If we consider an area where life starts from scratch by the process of succession, it is known as primary succession. However, if life starts at a place after the area has lost all the life forms existing there, the process is called secondary succession.

It is obvious that primary succession is a rather slow process as life has to start from nothing whereas secondary succession is faster because it starts at a place which had already supported life before. Moreover, the first species that comes into existence during primary succession is known as pioneer species.

Types of Ecological Succession

These are the following types of ecological succession:

Primary Succession

Primary succession is the succession that starts in lifeless areas such as the regions devoid of soil or the areas where the soil is unable to sustain life.

When the planet was first formed there was no soil on earth. The earth was only made up of rocks. These rocks were broken down by microorganisms and eroded to form soil. The soil then becomes the foundation of plant life. These plants help in the survival of different animals and progress from primary succession to the climax community.

If this primary ecosystem is destroyed, secondary succession takes place.

Secondary Succession

Secondary succession occurs when the primary ecosystem gets destroyed. For eg., a climax community gets destroyed by fire. It gets recolonized after the destruction. This is known as secondary ecological succession. Small plants emerge first, followed by larger plants. The tall trees block the sunlight and change the structure of the organisms below the canopy. Finally, the climax community arrives.