



RAMA
UNIVERSITY

www.ramauniversity.ac.in

FACULTY OF ENGINEERING & TECHNOLOGY
DEPARTMENT OF BIOTECHNOLOGY

LIST OF ENZYMES USED IN GENETIC ENGINEERING

1. Restriction Enzymes (Already discussed in the previous slides)
2. DNA Ligase (Already discussed in the previous slides)
3. DNA Polymerase I
4. Klenow Fragment
5. T₄ DNA Polymerase
6. Thermostable DNA Polymerases
7. Terminal Deoxinucleotidyl transferase
8. Reverse Transcriptase

DNA Polymerase I

The source of DNA polymerase I is *E.coli*

Function of DNA polymerase

- ❖ Exonuclease activity 5'-3'
- ❖ Exonuclease activity 3'-5'
- ❖ Polymerase activity 5'-3': addition of dNTP's at 3'-OH termini.
- ❖ Fills gaps in ds DNA



KLENOW FRAGMENT

- ❖ Klenow Fragment can be produced by treating with protease subtilisin.

Function of Klenow Fragment:


- ❖ Klenow fragment doesn't have 5' -3' Exonuclease activity. But has 3'-5' Exonuclease activity, and 5'-3' Polymerase activity.

LIGASE ENZYME

- The term Ligase is taken from the Latin verb ligāre, meaning of Ligare is "to bind" or to glue together.
- Ligase is an enzyme that can catalyze the joining of two large molecules by forming a new chemical bond, by hydrolysis of a small chemical group dependent to one of the larger molecules or the enzyme catalyzing the linking together of two compounds.
- In general, a ligase enzyme catalyzes the reaction:
$$Ab + C \rightarrow A-C + b$$

or

$$Ab + cD \rightarrow A-D + b + c \dots\dots$$
- Ligase can join two complementary fragments of nucleic acid by catalyzing the formation of a phosphodiester bond .
- DNA Ligase play role in
 1. DNA replication
 2. DNA repair



MECHANISM:

- Adenylation (addition of AMP) of a lysine residue in the active center of the enzyme takes place and pyrophosphate is released.
- Transfer of AMP to the 5' phosphate and pyrophosphate bond formation takes place.
- Formation of a phosphodiester bond between the 5' phosphate of the donor and the 3' hydroxyl of the acceptor.


LIST OF ENZYMES USED IN GENETIC ENGINEERING

T4 DNA Polymerase

T4 bacteriophage is the source of T4 DNA Polymerase.

Functions of T4 DNA Polymerase:

- ❖ It has 3'-5' exonuclease activity.
- ❖ It has 5'-3' Polymerase activity.
- ❖ It catalyzes template directed DNA synthesis.
- ❖ It has processivity (400 nucleotides/second).



Thermostable DNA Polymerases:

Thermostable DNA polymerase is usually isolated from thermophilic bacteria (like *Thermus aquaticus*).

Functions:

- ❖ The function of thermostable DNA polymerase is to catalyze template directed synthesis from free 3'-OH and bound to primer.
- ❖ It synthesizes DNA at high temperature (in case of PCR).
- ❖ For example: *Taq* DNA Polymerase, *pfu* DNA polymerase.


LIST OF ENZYMES USED IN GENETIC ENGINEERING

Terminal Deoxynucleotidyl Transferase:

Terminal deoxynucleotidyl transferase enzymes isolated from Immature, pre-B, pre-T lymphoid cells and acute lymphoblastic leukemia/lymphoma cells and commercially available terminal deoxynucleotidyl transferase enzyme is purified from recombinant *E. coli* cells expressing calf / rat / mouse thymus gene.

Functions:

- ❖ It adds a specific nucleotide to the 3'-end of a DNA strand
- ❖ It does not require a template
- ❖ It requires protruding 3' overhang



Reverse Transcriptase:

The source of Reverse transcriptase is retrovirus.

- For example:
1. Moloney murine leukemia virus (Mo-MLV)
 2. Avian myeloblastosis virus (AMV)

Functions:

Reverse Transcriptase enzymes have two types of activities

1. DNA Polymerase Activity
2. RNase H activity