



RAMA
UNIVERSITY

www.ramauniversity.ac.in

FACULTY OF NURSING



M.S.N 1ST UNIT-8TH DIABETES INCIPIDUS

RADHA DEVI
NURSING TUTOR
M.S.N DEPARTMENT
RAMA UNIVERSITY

DIABETES INCIPIDUS

Definition: It is a endocrine disorder ADH are affect the function of reabsorption of water and cause polyurea.

TYPES: 1.Craniogenic diabetes insipidus (ADH dependent)

2.Nephrogenic diabetes insipidus (Non ADH dependent)

CAUSES OF CRANIOGENIC DIABETES INCIPIDUS

- Brain Injury
- Posterior pituitary necrosis
- Hemorrhage
- Hypovolemic shock
- Decrease production of ADH

CAUSES OF NEPHROGENIC DIABETES INCIPIDUS

- Receptors problem
- Genetical causes
- Renal failure
- Drugs lithium

CLINICAL MANIFESTATIONS

- Polyurea
- Polydipcia
- Increase sodium level
- Dehydration
- Low B.P
- Lethargic and weakness
- Increase pulse rate in early stage

DIAGNOSIS

- C.T. Scan
- M.R.I.
- Asses ADH level
- Blood test

MEDICAL MANAGEMENT

- Craniogenic diabetes insipidus: provide ADH
 - Nephrogenic diabetes insipidus: Desmopressin
-
- **Nursing management:** To maintain hydration
 - Assess vital signs
 - Maintain input and output chart
 - Provide I.V. Fluid