


Types of culture media used in microbiology

Media are of different types on consistency and chemical composition.



A. On Consistency:

1. Solid Media. Advantages of solid media: (a) Bacteria may be identified by studying the colony character, (b) Mixed bacteria can be separated. Solid media is used for the isolation of bacteria as pure culture. 'Agar' is most commonly used to prepare solid media. Agar is polysaccharide extract obtained from seaweed. Agar is an ideal solidifying agent as it is : (a) Bacteriologically inert, i.e. no influence on bacterial growth,
- 2 (b) It remains solid at 37°C, and (c) It is transparent.
- 3 Liquid Media. It is used for profuse growth, e.g. blood culture in liquid media. Mixed organisms cannot be separated.

B. On Chemical Composition :

1. Routine Laboratory Media
2. Synthetic Media. These are chemically defined media prepared from pure chemical substances. It is used in research work.

ROUTINE LABORATORY MEDIA

These are classified into six types:

- (1) Basal media,
- (2) Enriched media,
- (3) Selective
- (4) Indicator media,
- (5) Transport media, and (6) Storage media.

1. **BASAL MEDIA.** Basal media are those that may be used for growth (culture) of bacteria that do not need enrichment of the media. Examples: Nutrient broth, nutrient agar and peptone water. Staphylococcus and Enterobacteriaceae grow in these media.

2. **ENRICHED MEDIA.** The media are enriched usually by adding blood, serum or egg. Examples: Enriched media are blood agar and Lowenstein-Jensen media. Streptococci grow in blood agar media.

3. **SELECTIVE MEDIA.** These media favour the growth of a particular bacterium by inhibiting the growth of undesired bacteria and allowing growth of desirable bacteria. Examples: MacConkey agar, Lowenstein-Jensen media, tellurite media (Tellurite inhibits the growth of most of the throat organisms except diphtheria bacilli). Antibiotic may be added

4. to a medium for inhibition.

5. **INDICATOR (DIFFERENTIAL) MEDIA.** An indicator is included in the medium. A particular organism causes change in the indicator, e.g. blood, neutral red, tellurite. Examples: Blood agar and MacConkey agar are indicator media.

6. **TRANSPORT MEDIA.** These media are used when specie-men cannot be cultured soon after collection. Examples: Cary-Blair medium, Amies medium, Stuart medium.

7. **STORAGE MEDIA.** Media used for storing the bacteria for a long period of time. Examples: Egg saline medium, chalk cooked meat broth

COMMON MEDIA IN ROUTINE USE

Nutrient Broth. 500 g meat, e.g. ox heart is minced and mixed with 1 litre water. 10 g peptone and 5 g sodium chloride are added, pH is adjusted to 7.3. Uses: (1) As a basal media for the preparation of other media, (2) To study soluble products of bacteria.

Nutrient Agar. It is solid at 37°C. 2.5% agar is added in nutrient broth. It is heated at 100°C to melt the agar and then cooled.

Peptone Water. Peptone 1% and sodium chloride 0.5%. It is used as base for sugar media and to test indole formation.

Blood Agar. Most commonly used medium. 5- 10% defibrinated sheep or horse blood is added to melted agar at 45-50°C. Blood acts as an enrichment material and also as an indicator. Certain bacteria when grown in blood agar produce haemolysis around their colonies. Certain bacteria produce no haemolysis. Types of changes : (a) beta (p) haemolysis. The colony is surrounded by a clear zone of complete haemolysis, e.g. *Streptococcus pyogenes* is a beta haemolytic streptococci, (b) Alpha (a) haemolysis. The colony is surrounded by a zone of greenish discolouration due to formation of biliverdin, e.g. *Viridans streptococci*, (c) Gamma (y) haemolysis, or, No haemolysis. There is no change in the medium surrounding the colony,

Chocolate Agar or Heated Blood agar:

Prepared by heating blood agar. It is used for culture of pneumococcus, gonococcus, meningo- coccus and *Haemophilus*. Heating the blood inactivates inhibitor of growths.

MacConkey Agar. Most commonly used for enterobac-teriaceae. It contains agar, peptone, sodium chloride, bile salt, lactose and neutral red. It is a selective and indicator medium :

(1) Selective as bile salt does not inhibit the growth of enterobacteriaceae but inhibits growth of many other bacteria.

(2) Indicator medium as the colonies of bacteria that ferment lactose take a pink colour due to production of acid. Acid turns the indicator neutral red to pink. These bacteria are called 'lactose fermenter', e.g. *Escherichia coll.* Colourless colony indicates that lactose is not

(3) fermented, i.e. the bacterium is non-lactose fermenter, e.g. *Salmonella*, *Shigella*, *Vibrio*.

(4)

(5) **Mueller Hinton Agar.** Disc diffusion sensitivity tests for antimicrobial drugs should be carried out on this media as per WHO recommendation to promote reproducibility and comparability of results.

(6)

(7) **Hiss's Serum Water Medium.** This medium is used to study the fermentation reactions of bacteria which can not grow in peptone water sugar media, e.g. pneumococcus, *Neisseria*, *Corynebacterium*.

(8)

(9) **Lowenstein-Jensen Medium.** It is used to culture tubercle bacilli. It contains egg, malachite green and glycerol. (1) Egg is an enrichment material which stimulates the growth of tubercle bacilli, (2) Malachite green inhibits growth of organisms other than mycobacteria, (3) Glycerol promotes the growth of *Mycobacterium tuberculosis* but not *Mycobacterium bovis*.

(10) **Dubos Medium.** This liquid medium is used for tubercle bacilli. In this medium drug sensitivity of tubercle bacilli can be carried out. It contains 'tween 80', bovine serum albumin, casein hydrolysate, asparagin and salts. Tween 80 causes dispersed growth and bovine albumin causes rapid growth.

(11) **Loeffler Serum.** Serum is used for enrichment. Diphtheria bacilli grow in this medium in 6 hours when the secondary bacteria do not grow. It is used for rapid diagnosis of diphtheria and to demonstrate volutin granules. It contains sheep, ox or horse serum.

(12) **Tellurite Blood Agar.** It is used as a selective medium for isolation of *Corynebacterium diphtheriae*

